

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

<sup>Rx</sup> Moxifloxacin & Dexamethasone Ophthalmic Solution

**LUMIMOX-D™**

Eye Drops

लुमिमोक्स-डी

**COMPOSITION :**

Moxifloxacin Hydrochloride	IP	
Equivalent to Moxifloxacin		0.5% w/v
Dexamethasone Sodium Phosphate	IP	0.1% w/v
Aqueous Buffered Vehicle		q.s.

**Dosage:** As directed by Physician

**CATEGORY:** Antibiotic with Steroid

**DESCRIPTION:**

**LUMIMOX-D** is a combination of Moxifloxacin 0.5% and Dexamethasone 0.1%.

Moxifloxacin is a fourth-generation fluoroquinolone, controls infection by inhibiting the DNAGyrase & topoisomerase iv. Dexamethasone is a potent corticosteroid effectively controls the inflammation by inhibiting the release of inflammatory mediators.

**INDICATIONS:**

**LUMIMOX-D** is indicated for steroid-responsive inflammatory ocular conditions for which a corticosteroid is indicated and where bacterial infection or a risk of bacterial ocular infection exists. The combination can also be used for post-operative inflammation and any other ocular inflammation associated with infection.

**CONTRA-INDICATIONS:**

**LUMIMOX-D** is contraindicated in epithelial herpes simplex keratitis (Dendritic keratitis), vaccinia, varicella, and in many other viral diseases of the conjunctiva and cornea, Mycobacterial infection of the eye and fungal diseases of ocular structures and in individuals hypersensitive to any of the components of the medication.

**Storage:** Store in cool & dark place.

**WARNINGS:**

- Remove contact lenses before using this eye drop.
- Use the solution within one month after opening the container.
- Do not use if the solution changes color or becomes cloudy.
- If eye pain, changes in vision, continued redness or irritation of the eye is experienced, or if the condition worsens or persists for more than 72 hours discontinue use and consult a medical doctor.

**PRECAUTIONS:**

Prolonged use of steroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision, and posterior subcapsular cataract formation.

**Adverse Reactions**

The most frequently reported drug-related undesirable effects seen with Moxifloxacin are conjunctival irritation, increased lacrimation, keratitis and papillary conjunctivitis. Secondary Infection.

**DOSAGE AND ADMINISTRATION:**

One or two drops instilled into the conjunctival sac(s), every 4 to 6 hours. During the initial 24 to 48 hours, the dosage may be increased to 1 or 2 drops every two hours. Frequency must be decreased gradually or warranted by improvement in clinical signs. Care should be taken not to discontinue the therapy prematurely. Or as directed by the Physician.

Keep out of reach of children.

For external use only.

Not for injection.

**PRESENTATION:**

**LUMIMOX-D** is a sterile solution supplied in white opaque plastic dropper bottle with a cap, containing 5ml of the solution.

**Mfg. Lic. No. G/28/1536**

Manufactured in India by :



Marketed by:



**LUMIN LABS LLP**

D2-304, Siyona Pethkar, Punawale,  
Mumbai - Bangalore Highway,  
Pune-411033

50 X 100 MM